

A search for cool subdwarfs: Stellar parameters for 134 candidates¹

David Yong and David L. Lambert

Department of Astronomy, University of Texas, Austin, TX 78712

tofu,dll@astro.as.utexas.edu

ABSTRACT

The results of a search for cool subdwarfs are presented. Kinematic (U, V, and W) and stellar parameters (T_{eff} , $\log g$, $[\text{Fe}/\text{H}]$, and ξ_t) are derived for 134 candidate subdwarfs based on high resolution spectra. The observed stars span $4200\text{K} < T_{\text{eff}} < 6400\text{K}$ and $-2.70 < [\text{Fe}/\text{H}] < 0.25$ including only 8 giants ($\log g < 4.0$). Of the sample, 100 stars have MgH bands present in their spectra. The targets were selected by their large reduced proper-motion, the offset from the solar metallicity main sequence, or culled from the literature. We confirm the claims made by Ryan (1989) regarding the NLTT catalog being a rich source of subdwarfs and verify the success of the reduced proper-motion constraint in identifying metal-poor stars.

Subject headings: stars: abundances – stars: fundamental parameters – subdwarfs

1. Introduction

The driving force behind our understanding of the chemical evolution of the Galaxy is the interpretation of observed abundance ratios. Self-consistent analyses of large high quality data sets have revealed detailed abundance patterns (e.g., Edvardsson et al. 1993, McWilliam et al. 1995). Recent attempts to understand the observed abundance trends include endeavors by Timmes et al. (1995), Goswami & Prantzos (2000), and Alibés et al. (2001) who predict the evolution of the abundances of all elements from carbon through zinc. Cool stars provide a unique opportunity to test directly these models of Galactic chemical

¹Based in-part on observations obtained with the Hobby-Eberly Telescope, which is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig-Maximilians-Universität München, and Georg-August-Universität Göttingen.

evolution through the abundances of low ionization potential trace elements (e.g., Rb, Cs) and isotopic ratios measured from molecular bands (e.g., Mg from MgH, Ti from TiO). Presently these tests cannot be carried out due to the dearth of known cool metal-poor stars. In order to rectify this situation we have commenced a search for cool subdwarfs, unevolved metal-poor stars that fall below the solar metallicity main sequence in color-magnitude diagrams.

Searches undertaken by Carney & Latham (1987) and Ryan (1989) selected candidates drawn from proper-motion catalogs. These searches were successful in identifying metal-poor dwarfs for subsequent detailed abundance analysis. However, both searches neglected cool subdwarfs. An alternative to searches for metal-poor stars based on proper-motion catalogues are the objective prism surveys which identify candidate metal-poor stars by the weakening of Ca II H and K. Such studies have been conducted by Bond (1970), Bond (1980), Beers et al. (1985), Beers et al. (1992), and others. Selection criteria which rely exclusively upon Ca features as metallicity indicators strongly bias the sample towards warmer stars. Since metal lines weaken with increasing T_{eff} , a temperature estimate is required before deciding if lines are abnormally weak. In the absence of temperature indicators, a cool metal-poor star will have Ca II features comparable to a warmer solar metallicity star. More current work by Christlieb (2000) on the stellar content of the Hamburg/ESO Survey has dramatically increased the yields of metal-poor stars. Candidate metal-poor stars are identified from their location in $(B - V) - \text{Ca line strength}$ space. By comparing the Ca line strength between stars of similar T_{eff} 's, this increases the likelihood that a metal deficiency is the cause of the relative weakening of the Ca line in a candidate. With the remarkable success rate of 60% for stars below $[\text{Fe}/\text{H}] = -2.0$, twice as good as the HK survey by Beers and collaborators, we look forward to the results of the Hamburg/ESO Survey when applied to cool subdwarfs.

In this paper we present the results of our search for cool subdwarfs, stellar (T_{eff} , $\log g$, $[\text{Fe}/\text{H}]$, and ξ_t) and kinematic parameters (U, V, and W) for 134 stars. Our sample includes 80 stars with no prior metallicity estimates. In §2 we outline and justify the selection criteria. The observations will be described in §3 and the analysis in §4. A discussion will be presented in §5 and concluding remarks given in §6.

2. Selection criteria

Our goal was to find previously unidentified subdwarfs with $4000\text{K} < T_{\text{eff}} < 4700\text{K}$ and $[\text{Fe}/\text{H}] < -1.5$. We modelled our search upon the highly successful effort by Ryan (1989) using the NLTT catalog of stars with annual proper-motions in excess of $0''.18/\text{year}$ (Luyten 1979, 1980; Luyten & Hughes 1980). Subdwarfs are on plunging orbits, often highly elliptical,

resulting in large transverse velocities relative to the local standard of rest. Subdwarfs are therefore over-represented in proper-motion catalogues. Following Ryan (1989), our primary criterion for ensuring that metal-poor candidates were selected was the reduced proper-motion, $H = m_V + 5 \log \mu + 5$ where m_V is the apparent magnitude and μ is the proper-motion in arcsec/year. The reduced proper-motion can also be expressed as $H = M_V + 5 \log \nu_T - 3.37$ where M_V is the absolute magnitude and ν_T is the transverse velocity in km/s. At a given spectral type, a reduced proper-motion constraint rejects stars whose transverse velocities fall below a chosen value. Subdwarfs are less luminous than disk dwarfs at a given color and have larger transverse velocities. Both effects conspire to increase the reduced proper-motion of the subdwarf population relative to the disk dwarf population. We adopted Ryan’s reduced proper-motion constraint requiring stars have $H_R \geq 10.7 + 2(m_{pg} - m_R)$ where $H_R = m_R + 5 \log \mu + 5$ using Luyten magnitudes m_R and m_{pg} . We imposed a magnitude limit $m_R \leq 13.0$ and a declination limit $-40^\circ \leq \delta \leq 90^\circ$ as our observations were made at McDonald Observatory where we required reasonable signal-to-noise (60 per pixel at 6500Å) in 20-30 minute exposures. To ensure that we observed cool stars, Luyten color classes g-k, k, and k-m were selected. Carney & Latham (1987) and Ryan (1989) previously observed the g-k color class. We included these stars so that we could compare our derived stellar parameters with the Carney and Ryan values. Ryan’s photometry suggested that the color classes were not accurate. Stars assigned color class g-k by Luyten actually span late F to late K. Further, the effect of the ultraviolet excess, the hallmark of a subdwarf, is to camouflage a cool subdwarf as a warmer disk dwarf in the Luyten color system. Altogether, our selection criteria resulted in 4445 NLTT candidate subdwarfs.

Our list was augmented by subdwarfs previously identified in the literature. Our final sources of candidate subdwarfs were color-magnitude diagrams constructed using the HIPPARCOS (ESA 1997) and Yale (van Altena et al. 1995) parallaxes. Targets located beneath the solar metallicity main sequence in the range $0.8 < B - V < 1.4$ were selected. Subdwarfs have a higher T_{eff} than disk dwarfs at a given mass which shifts subdwarfs blueward of the disk main sequence. Reid et al. (2001) demonstrated that in the range $B - V < 0.8$, the majority of stars below the disk main sequence in a color-magnitude diagram based on HIPPARCOS data are not subdwarfs, $[\text{Fe}/\text{H}] > -1.0$. Errors in the colors cause these disk dwarfs to appear as subdwarfs. Our observations and analysis will test if the Reid et al. (2001) findings are valid when we extend to cooler dwarfs, $0.8 < B - V < 1.4$. In total, we observed around 230 candidate subdwarfs though in this paper we are reporting on 134 candidates.

3. Observations and data reduction

Table 1 contains the list of candidates observed at McDonald Observatory on the 2.7m Harlan J. Smith telescope and on the 9.2m Hobby-Eberly telescope (HET) between November 1999 and April 2002. The 2.7m data were obtained using the cross-dispersed echelle spectrometer (Tull et al. 1995) at the coudé f/32.5 focus with a resolving power of either 30,000 or 60,000. The detector was a Tektronix CCD with $24 \mu\text{m}^2$ pixels in a 2048×2048 format. For this setting, the spectral coverage was from 3800Å to 8900Å with gaps between the orders beyond 5800Å. The HET data were taken with the Upgraded Fiber Optic Echelle spectrograph (Harlow et al. 1996) at a resolving power of 11,000 on a 1024×1024 CCD. The spectral coverage was from 4500Å to 9000Å with gaps between the orders beyond 7300Å. For observations on both telescopes, wavelength coverage incorporated the MgH A-X bands near 5140Å as well as the TiO (0,0) bandhead of the γ -system $\text{A}^3\Phi - \text{X}^3\Delta$ near 7054Å. Visual inspection of the spectra readily indicated the presence or absence of the MgH or TiO molecular features (see Figures 1 and 2). Of our sample of 134 stars, 34 displayed neither MgH nor TiO bands and 100 showed MgH or MgH and TiO bands. Numerous Fe I, Fe II, Ti I, and Ti II lines were available in the observed spectra for spectroscopic determination of stellar parameters. For each star, exposure times were generally 20-30 minutes and only occasionally were multiple exposures taken and co-added. Although varying from star to star, the typical signal-to-noise of the extracted one dimensional spectra was 60 per pixel at 6500Å. One dimensional wavelength calibrated normalized spectra were extracted in the standard way using the IRAF² package of programs. Equivalent widths were measured using IRAF where in general Gaussian profiles were fit to the observed profile. Figure 3 shows the spectra of 2 different stars, one taken with R=60,000 and the other at R=11,000 where two Fe I lines used in the analysis are indicated. Although less than ideal, an equivalent width analysis of R=11,000 spectra is feasible.

4. Analysis

4.1. Deriving stellar parameters

The LTE stellar line analysis program MOOG (Sneden 1973) was used in combination with the adopted model atmosphere. For $\log g > 3.5$, we used the NEXTGEN model atmo-

²IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

sphere grid for low mass stars computed by Hauschildt et al. (1999). For giants with $\log g \leq 3.5$ we used LTE model atmospheres computed by Kurucz (1993). In both cases we interpolated within the grid when necessary. We derived the model parameters in the following way. We measured the equivalent widths of ~ 35 Fe I and ~ 5 Fe II lines. Values for T_{eff} were set from excitation equilibrium. T_{eff} was adjusted so that Fe abundances derived from individual lines were independent of excitation potential, as they must be. The Fe abundances derived from individual Fe I lines must be independent of the strength of the line. Thus the microturbulence, ξ_t , was estimated. Finally, the gravity was determined from ionization equilibrium. That is, the Fe abundance derived from Fe I lines must agree with the abundance derived from Fe II lines. This process was iterated until a consistent set of parameters were obtained (see Figure 4). Ti I and Ti II lines were also used to check the stellar parameters. The final value of $[\text{Fe}/\text{H}]$ was simply the weighted mean of the Fe I and Fe II abundances derived from the accepted model (see Table 1 for model parameters).

The gf -values of the Fe lines used in the analysis were selected from Lambert et al. (1996) and a list compiled by R. E. Luck (1993, private communication). The gf -values of the Ti lines were selected from the R. E. Luck compilation. Where possible, only weak lines, $\text{EWs} < 90 \text{ m}\text{\AA}$, were used in the analysis. We checked our analysis techniques by observing the solar spectrum at $R=60,000$. We measured 30 Fe I lines and 7 Fe II lines and compared our equivalent widths with the Grevesse & Sauval (1999) values. Our equivalent widths were larger by a mean value of $3.7 \text{ m}\text{\AA}$ with a standard deviation of $2.6 \text{ m}\text{\AA}$. Using our equivalent widths and a NEXTGEN model atmosphere, we derived a solar abundance of $\log \epsilon (\text{Fe}) = 7.54$. Considering the Grevesse & Sauval (1999) value of $\log \epsilon (\text{Fe}) = 7.50 \pm 0.05$ derived from their empirical model solar atmosphere, we adopted $\log \epsilon (\text{Fe}) = 7.52$ as the solar Fe abundance for this study.

Our derived model parameters, T_{eff} , $\log g$, ξ_t , and $[\text{Fe}/\text{H}]$ have associated uncertainties. We varied T_{eff} until the trends between lower excitation potential and abundance were unacceptable. Similarly, we took values for ξ_t that noticeably changed the trends between equivalent width and abundance. For $\log g$, we measured the standard deviation of the Fe abundance as derived from Fe I lines (typically 0.1 to 0.15 dex) then allowed Fe I and Fe II to agree within this standard deviation. This produced uncertainties of $\delta T_{\text{eff}}=150\text{K}$, $\delta \log g=0.3 \text{ dex}$, $\delta \xi_t=0.3 \text{ km/s}$, and $\delta [\text{Fe}/\text{H}]=0.2 \text{ dex}$ in the model parameters.

4.2. Comparison with literature

For the 134 candidate subdwarfs analyzed here, a search on SIMBAD indicated that there were 37 stars with previously determined values for T_{eff} and 54 stars with previously

determined values for $[\text{Fe}/\text{H}]$. We have compared our values with those found in the literature (see Table 2 and Figures 5 and 6). We find a mean offset $\langle T_{\text{eff}}(\text{this study}) - T_{\text{eff}}(\text{literature}) \rangle = -25$ K with a standard deviation of 114K. For $[\text{Fe}/\text{H}]$, the mean offset is $\langle [\text{Fe}/\text{H}] (\text{this study}) - [\text{Fe}/\text{H}] (\text{literature}) \rangle = -0.1$ dex with a standard deviation of 0.38 dex. The agreement is reasonable between the stellar parameters derived in this study and the values found in a variety of sources in the literature.

Table 2 shows values for $[\text{Fe}/\text{H}]$ given by Ryan & Norris (1991) for a number of stars. However, these metallicities are not paired with values for T_{eff} . The Ryan & Norris (1991) sample are re-observations of a subset of metal-weak stars identified in Ryan (1989). An index measuring the strength of the Ca II K line was calibrated empirically as a function of $B - V$ to give $[\text{Fe}/\text{H}]$. Alonso et al. (1996a, 1999a) determined T_{eff} 's using the infrared flux method which requires a metallicity and gravity estimate. The derived T_{eff} 's are insensitive to the input metallicity and gravity. We also computed T_{eff} 's derived by use of the $T_{\text{eff}}:[\text{Fe}/\text{H}]:\text{color}$ relations given in Alonso et al. (1996b, 1999b) assuming values for $[\text{Fe}/\text{H}]$ from this study, see Table 3.

MgH and TiO bands will not be visible in the spectra of stars with sufficiently high temperatures and/or low abundances. Indeed, this is the case for 34 of the stars as noted in the last column of Table 1.

5. Discussion

5.1. Kinematics

We selected stars with large reduced proper-motions and therefore expect the observed sample to be kinematically distinct from the thin disk. Following the prescription given by Johnson & Soderblom (1987), the Galactic space-velocity components U , V , and W were calculated along with the associated uncertainties (see Table 1). To correct for the solar motion with respect to the local standard of rest (LSR) we assumed the Dehnen & Binney (1998) values $(-10, +5, +7)$ km s $^{-1}$ in (U, V, W) . In the absence of HIPPARCOS parallaxes, spectroscopic parallaxes were determined by using the derived model parameters and the Girardi et al. (2000) isochrones. Figure 7 shows $[\text{Fe}/\text{H}]$ versus U_{LSR} , V_{LSR} , and W_{LSR} . We identify stars which lag the LSR, $V < -50$ km/s, as being members of the thick disk or halo. As expected from a reduced proper-motion constraint, we have selected stars which belong to populations kinematically distinct from the thin disk. Many of the observed stars have abundances indicative of the thick disk and halo. Ryan (1989) claimed that the NLTT catalog was a rich source of subdwarfs as identified by the ultraviolet excess. Figure 8 shows

the numbers of stars for various metallicity bins. Given that 27 stars have $[\text{Fe}/\text{H}] \leq -1.5$, our data endorse Ryan’s claims. From the kinematics and abundances of the observed stars, we conclude that the reduced proper-motion constraint successfully selected metal-weak thick disk and halo stars.

5.2. Temperature

Our goal was to find cool stars, $4000\text{K} < T_{\text{eff}} < 4700\text{K}$. Of the 134 stars presented in this paper, 44 had $T_{\text{eff}} \leq 4700\text{K}$ and given the uncertainties in T_{eff} we note that 69 of the 134 stars had $T_{\text{eff}} \leq 4800\text{K}$. Figure 9 shows the number of stars versus T_{eff} for this study and the Carney et al. (1994) study. Our sample includes cooler stars than the Carney study which highlights the different temperature regimes of interest. In addition to the stars presented here, we have observations of a further 100 stars which are simply too cool for equivalent width analysis. Molecular bands make identification of the continuum virtually impossible and finding unblended Fe I and Fe II lines is problematic. Given the strength of the TiO bands, we assume that these stars are all cooler than 4500K. We are exploring various techniques for determination of the stellar parameters. The strength of the TiO bandheads in some stars suggests rather low values for T_{eff} , thus we reconfirm Ryan’s finding regarding Luyten’s photometry that “the values tabulated in the NLTT catalog must be regarded as approximate only”.

5.3. Metallicity

Figure 8 shows the number of stars versus metallicity for this study, the Carney et al. (1994) sample, and the Ryan & Norris (1991) sample. It is clear that our sample is considerably smaller than the Ryan and Carney studies. The distributions are similar between the Carney sample and our study which is rather interesting. Our complete list of NLTT candidates comprised 4445 stars of which we observed some 230 stars. We derived stellar parameters for 134 stars and found 27 with $[\text{Fe}/\text{H}] \leq -1.50$, with several having prior metallicity estimates. Presumably, if we were to observe the remaining 4000+ candidates, we would find ~ 500 stars with $[\text{Fe}/\text{H}] \leq -1.50$. The distribution of the Ryan sample differs from this study and from the Carney study. The Ryan sample peaks at a lower metallicity where this is simply a selection effect. The original Ryan (1989) sample provided broadband photometry where metal-poor candidates were identified by the ultraviolet excess. The stars re-observed in Ryan & Norris (1991) were those stars identified in Ryan (1989) as having an ultraviolet excess corresponding to $[\text{Fe}/\text{H}] < -1.2$, that is, $\delta(U - B)_{0.6} > 0.2$.

For the 27 stars we observed with $[\text{Fe}/\text{H}] \leq -1.50$, there were 11 sufficiently cool to provide MgH features. These were the cool subdwarfs with molecular features that we were interested in finding. Again, if we were to observe the remaining 4000+ NLTT candidates there would presumably be ~ 200 such objects.

In our sample, the lack of cool subdwarfs with $[\text{Fe}/\text{H}] \leq -2$ could be attributed to 2 reasons. Firstly, the volume we are sampling may not be large enough and secondly, our ability to identify genuine subdwarfs may be inefficient. Regarding the volume we are sampling, according to the Girardi et al. (2000) isochrones, for stars with $T_{\text{eff}} = 4500\text{K}$ and $[\text{Fe}/\text{H}] = -1.68$, our $m_R \simeq V = 13$ limit corresponds to a maximum distance of merely ~ 75 parsecs. This volume further decreases when we consider even more metal-poor stars. More importantly, applying our magnitude limit to a star with $T_{\text{eff}} = 5500\text{K}$ and $[\text{Fe}/\text{H}] = -1.68$ corresponds to a maximum distance of ~ 220 parsecs. This represents a volume 27 times larger than for our cooler targets. Carney et al. (1994) considered stars as faint as $V=16$ which combined with the intrinsic brightness of warmer targets results in a considerably larger volume. Obviously the number of cool metal-poor stars in a particular volume increases as the volume increases. Regarding the efficiency of identifying candidate subdwarfs, recent work by Salim & Gould (2002) provides more accurate $V - J$ photometry for NLTT stars based on 2MASS infrared photometry and USNO-A optical photometry. Combining the precise photometry with the proper-motions generates a reduced proper-motion diagram with well defined white dwarf, subdwarf, and disk dwarf populations. From this revised NLTT catalog and proper-motion diagram we expect that these subdwarf candidates will contain a considerably higher fraction of genuine subdwarfs than our original list of candidates. Observations of candidate subdwarfs selected from the Salim & Gould (2002) $V - J$ reduced proper-motion diagram will be made.

Alternatively, we could consider what volume is needed to obtain a large number of stars with $-3 < [\text{Fe}/\text{H}] < -2$ and $4000\text{K} < T_{\text{eff}} < 4500\text{K}$. Our observed NLTT candidates provided just 1 star within the desired range of parameters, G 39-36 ($T_{\text{eff}}=4200\text{K}$, $[\text{Fe}/\text{H}]=-2.5$). From the Yi et al. (2001) isochrones, we estimate that G 39-36 lies at a distance of 27 parsecs from the sun. Given the stellar parameters of G 39-36, our $V=13$ magnitude limit represents a maximum volume of merely 36 parsecs. Since we observed around 200 of the 4445 NLTT candidates, perhaps there are 10 to 20 cool subdwarfs amongst our list of NLTT candidates. If we conservatively estimate that there are 5 cool subdwarfs ($-3 < [\text{Fe}/\text{H}] < -2$, $4000\text{K} < T_{\text{eff}} < 4500\text{K}$) within a radius of 50 parsecs around the sun, then we expect around 300 cool subdwarfs within a radius of 200 parsecs. A search for subdwarfs similar to G 39-36 out to a distance of 200 parsecs would require a magnitude limit of $V \simeq 17$. Such a limit implies a low resolution, low signal-to-noise search technique or a photometric search to weed out the intruders.

5.4. Hipparcos

Having observed a number of cool stars contained in the HIPPARCOS catalog, we can comment upon the results of Reid et al. (2001). The majority of HIPPARCOS stars were selected by their large reduced proper-motion from the NLTT catalog rather than the position in the color-magnitude diagram. Reid et al. (2001) claimed that in a color-magnitude diagram produced from HIPPARCOS parallaxes, the majority of stars with $(B - V) \leq 0.8$ situated below the solar metallicity main sequence were incorrectly positioned due to errors in the colors. These stars, located where subdwarfs ought to lie, were metal rich disk dwarfs rather than subdwarfs. Reid et al. (2001) also showed that the Tycho colors $(B - V)_T$ are redder than the Johnson $(B - V)$ colors. In Figure 10, we take the Tycho colors and HIPPARCOS parallaxes to produce a color-magnitude diagram overlaying the stars we observed. Nearby stars ($\pi > 32$ mas) with accurate parallaxes ($\sigma_\pi/\pi < 0.15$) are included to represent the disk population, $[\text{Fe}/\text{H}] \simeq 0$. Our few observations confirm and extend the Reid et al. (2001) findings in the regime $0.8 < B - V < 1.4$. Metal-poor stars with $[\text{Fe}/\text{H}] < -1.0$ are located amongst disk stars. Also, metal-rich stars with $[\text{Fe}/\text{H}] > -1.0$ are offset from the disk main sequence. The effect of the redder Tycho colors is evident as the disk main sequence is systematically offset from the Girardi et al. (2000) $[\text{Fe}/\text{H}] = 0$ isochrone. We intend to observe additional HIPPARCOS subdwarf candidates in the range $0.8 < (B - V)_T < 1.2$ not only to find cool subdwarfs but to further test the findings made by Reid et al. (2001).

6. Concluding remarks

We present stellar parameters for 134 candidate subdwarfs selected by their reduced proper-motion, offset from the solar metallicity, or from the literature. Our goal was to provide a large database of cool subdwarfs. Our selection criteria were successful in identifying cool stars (69 with $T_{\text{eff}} < 4800\text{K}$) and metal-poor stars (27 with $[\text{Fe}/\text{H}] \leq -1.50$). Of our sample, 11 stars were sufficiently cool to provide measurable MgH lines with $[\text{Fe}/\text{H}] < -1.50$. Armed with a sample of cool subdwarfs, we can begin to exploit their unique qualities. For a subset of our sample, we have measured the Mg isotopic abundance ratios and compared the observed trends with predictions from models of Galactic chemical evolution. This work will be presented in a future paper. We also intend to make further observations of candidate subdwarfs. Targets will be selected from the revised NLTT catalog (Salim & Gould 2002). Cool HIPPARCOS subdwarf candidates will be observed and we anticipate that Christlieb and collaborators will identify cool subdwarfs in the Hamburg/ESO survey data.

We thank George Preston for helpful comments and the staff of the Hobby-Eberly

telescope for service observations of our targets. We acknowledge support via the grant F-634 from the Robert A. Welch Foundation of Houston, Texas. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France and NASA’s Astrophysics Data System.

REFERENCES

- Alibés, A., Labay, J., & Canal, R. 2001, *A&A*, 370, 1103
- Alonso, A., Arribas, S., & Martínez-Roger, C. 1996a, *A&AS*, 117, 227
- . 1996b, *A&A*, 313, 873
- . 1999a, *A&AS*, 139, 335
- . 1999b, *A&AS*, 140, 261
- Beers, T. C., Preston, G. W., & Shectman, S. A. 1985, *AJ*, 90, 2089
- . 1992, *AJ*, 103, 1987
- Beers, T. C., Rossi, S., Norris, J. E., Ryan, S. G., & Shefler, T. 1999, *AJ*, 117, 981
- Bond, H. E. 1970, *ApJS*, 22, 117
- . 1980, *ApJS*, 44, 517
- Carney, B. W. & Latham, D. W. 1987, *AJ*, 93, 116
- Carney, B. W., Latham, D. W., Laird, J. B., & Aguilar, L. A. 1994, *AJ*, 107, 2240
- Carney, B. W., Wright, J. S., Sneden, C., Laird, J. B., Aguilar, L. A., & Latham, D. W. 1997, *AJ*, 114, 363
- Christlieb, N. 2000, Ph.D. Thesis
- Clementini, G., Gratton, R. G., Carretta, E., & Sneden, C. 1999, *MNRAS*, 302, 22
- Dehnen, W. & Binney, J. J. 1998, *MNRAS*, 298, 387
- Edvardsson, B., Andersen, J., Gustafsson, B., Lambert, D. L., Nissen, P. E., & Tomkin, J. 1993, *A&A*, 275, 101
- Eggen, O. J. 1987, *AJ*, 93, 393

- . 1998, *AJ*, 115, 2397
- ESA. 1997, *VizieR Online Data Catalog*, 1239
- Fuchs, B. & Jahreiß, H. 1998, *A&A*, 329, 81
- Fulbright, J. P. 2000, *AJ*, 120, 1841
- Gay, P. L. & Lambert, D. L. 2000, *ApJ*, 533, 260
- Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, *A&AS*, 141, 371
- Gizis, J. E. 1997, *AJ*, 113, 806
- Goswami, A. & Prantzos, N. 2000, *A&A*, 359, 191
- Grevesse, N. & Sauval, A. J. 1999, *A&A*, 347, 348
- Harlow, J. J. B., Ramsey, L. W., Andersen, D. R., Fleig, J. D., Rhoads, B. T., & Engel, L. G. 1996, *BAAS*, 28, 1324
- Hauschildt, P. H., Allard, F., & Baron, E. 1999, *ApJ*, 512, 377
- Johnson, D. R. H. & Soderblom, D. R. 1987, *AJ*, 93, 864
- Jorgensen, U. G. 1994, *A&A*, 284, 179
- Kurucz, R. 1993, *ATLAS9 Stellar Atmosphere Programs and 2 km/s grid*. Kurucz CD-ROM No. 13. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1993., 13
- Lambert, D. L., Heath, J. E., Lemke, M., & Drake, J. 1996, *ApJS*, 103, 183
- Luyten, W. J. 1979,1980, *New Luyten Catalogue of Stars with Proper Motions Larger than Two Tenths of an Arcsecond* (Minneapolis: University of Minnesota, 1979,1980)
- Luyten, W. J. & Hughes, H. S. 1980, *Proper Motion Survey with the Forty-Eight Inch Schmidt Telescope. LV. First Supplement to the NLTT Catalogue* (Minneapolis: University of Minnesota, 1980)
- McWilliam, A., Preston, G. W., Sneden, C., & Searle, L. 1995, *AJ*, 109, 2757
- Morrison, H. L., Olszewski, E. W., Mateo, M., Norris, J. E., Harding, P., Dohm-Palmer, R. C., & Freeman, K. C. 2001, *AJ*, 121, 283
- Norris, J. 1986, *ApJS*, 61, 667

- Primas, F., Molaro, P., Bonifacio, P., & Hill, V. 2000, *A&A*, 362, 666
- Reid, I. N., van Wyk, F., Marang, F., Roberts, G., Kilkenny, D., & Mahoney, S. 2001, *MNRAS*, 325, 931
- Ryan, S. G. 1989, *AJ*, 98, 1693
- Ryan, S. G. & Norris, J. E. 1991, *AJ*, 101, 1835
- Salim, S. & Gould, A. 2002, *ApJ*, 575, L83
- Schuster, W. J., Parrao, L., & Contreras Martinez, M. E. 1993, *A&AS*, 97, 951
- Skrutskie, M. F., Schneider, S. E., Stiening, R., Strom, S. E., Weinberg, M. D., Beichman, C., Chester, T., Cutri, R., Lonsdale, C., Elias, J., Elston, R., Capps, R., Carpenter, J., Huchra, J., Liebert, J., Monet, D., Price, S., & Seitzer, P. 1997, in *ASSL Vol. 210: The Impact of Large Scale Near-IR Sky Surveys*, 25
- Snedden, C. A. 1973, Ph.D. Thesis
- Thévenin, F. & Idiart, T. P. 1999, *ApJ*, 521, 753
- Timmes, F. X., Woosley, S. E., & Weaver, T. A. 1995, *ApJS*, 98, 617
- Tomkin, J. & Lambert, D. L. 1999, *ApJ*, 523, 234
- Tull, R. G., MacQueen, P. J., Sneden, C., & Lambert, D. L. 1995, *PASP*, 107, 251
- van Altena, W. F., Lee, J. T., & Hoffleit, D. 1995, *VizieR Online Data Catalog*, 1174
- Yi, S., Demarque, P., Kim, Y., Lee, Y., Ree, C. H., Lejeune, T., & Barnes, S. 2001, *ApJS*, 136, 417




Fig. 1.— $R=60,000$ data showing the absence (upper) and presence (lower) of the MgH A-X lines near $5140\text{\AA}$. Many of the additional lines in the lower spectrum are due to MgH.




Fig. 2.— R=60,000 data showing the absence (upper) and presence (lower) of the TiO γ -system $A^3\Phi-X^3\Delta$ bandhead near 7054Å.




Fig. 3.— Spectra of G 103-27 taken at $R=60,000$ and G 9-31 take at $R=11,000$. Both stars have similar stellar parameters. Representative Fe I lines used to derive stellar parameters are highlighted. Importantly, the $R=11,000$ spectra has Fe I lines for which equivalent widths can be measured.




Fig. 4.— Determination of stellar parameters T_{eff} , $\log g$, and ξ_t using excitation and ionization equilibrium for GJ 1064B. In the top panel, the lower excitation potential (LEP)-abundance relation is used to set T_{eff} . In the middle panel, the reduced equivalent width (W_λ/λ)-abundance relation is used to determine ξ_t . In the bottom panel, the abundances of Fe I (crosses) and Fe II (filled circles) are used to fix $\log g$. In all panels the line represents the least squares fit to the data.




Fig. 5.— T_{eff} comparisons between this work and various literature studies. The filled circles represent stars which have neither MgH nor TiO bands in their spectra. The dotted line represents the line of equality.




Fig. 6.— $[\text{Fe}/\text{H}]$ comparisons between this work and various literature studies. The filled circles represent stars which have neither MgH nor TiO bands in their spectra. The dotted line represents the line of equality. We note that the scatter for the open circles is slightly larger than for the filled circles.




Fig. 7.— Galactic space-velocity U, V , and W versus $[\text{Fe}/\text{H}]$ where U, V , and W are relative to the local standard of rest (LSR). Filled circles represent stars which have neither MgH nor TiO bands in their spectra. Only stars which have $\frac{\sigma_U + \sigma_V + \sigma_W}{|U| + |V| + |W|} < 0.7$ are shown, 117 out of 134. A considerable fraction of the sample noticeably lag the LSR ($V < -50$ km/s)




Fig. 8.— Number of stars versus metallicity for this study (upper), the Carney et al. (1994) study (middle), and the Ryan & Norris (1991) study (lower). Note the similar distributions between this study and the Carney work. The Ryan distribution peaks at a lower metallicity since their sample consists of stars identified as having an ultraviolet excess corresponding to $[\text{Fe}/\text{H}] < -1.2$.




Fig. 9.— Number of stars versus T_{eff} for this study (upper) and the Carney et al. (1994) study (lower). In the upper panel, the filled in histogram represents the distribution of stars with $[\text{Fe}/\text{H}] \leq -1.0$. The distribution for our sample peaks at lower values of T_{eff} than the Carney sample. This was expected since we deliberately selected cooler color classes.




Fig. 10.— The M_V , $(B - V)_T$ color-magnitude diagram with crosses marking HIPPARCOS stars with $\sigma_\pi/\pi < 15$ and $\pi > 32\text{mas}$. Closed circles represent observed stars with $[\text{Fe}/\text{H}] \geq -1.0$ while open circles represent $[\text{Fe}/\text{H}] < -1.0$. Girardi et al. (2000) isochrones using Johnson $(B - V)$ for $[\text{Fe}/\text{H}] = 0.0, -0.68, -1.68$ are overplotted. $(B - V)_T$ is increasingly redder than the Johnson $(B - V)$ as the stars get cooler as noted by Reid et al. (2001).

Table 1. Basic data and derived parameters for objects

Name	Ra ^a	Dec ^b	Resolving	T _{eff}	[Fe/H]	log g	ξ_t	U _{LSR}	σ_U	V _{LSR}	σ_V	W _{LSR}	σ_W	Luyten color	Molecular
	2000	2000	Power ^c	(K)		(cm s ⁻²)				(km/s)				or source	Features
PLX 5805	000155	260015	30000	4600	-1.72	4.25	1.0	120	10	-97	3	-26	8	Yale	Y
LP 524-64	000800	081642	30000	4600	-1.04	5.00	1.0	-90	30	-95	25	8	13	k-m	Y
BD -14 142	004822	-140713	60000	6200	-1.21	4.00	1.0	50	30	-225	110	2	25	g-k	N
LP 646-24	005006	-042600	30000	4800	-1.01	4.50	0.7	-16	5	-120	45	-38	16	g-k	Y
G 72-34	014604	355448	11000	4800	-2.23	4.75	0.9	45	10	-265	165	-243	230	g-k	N
BD -4 290	015218	-032643	60000	4900	-0.01	4.50	0.7	50	10	-2	1	-10	5	k	Y
BD +8 335	021120	093724	60000	5100	-0.98	4.50	0.7	-35	2	-65	13	-54	5	g-k	Y
HIP 10412	021411	013312	30000	4600	-0.40	4.75	1.2	-65	6	-24	5	2	3	Yale	Y
Cool 340	021726	270827	60000	4800	-0.40	5.00	0.8	14	6	-98	17	16	7	g-k	Y
BD +4 415	023435	052647	60000	4750	-0.63	4.50	0.6	122	6	-52	3	-15	6	k	Y
G 78-26	031627	380556	30000	4200	-1.01	4.75	0.6	65	5	-245	15	-17	6	Gizis97	Y
GJ 1064A	034702	412538	60000	5000	-1.15	4.50	0.6	-106	3	-109	7	-71	4	FJ98	Y
GJ 1064B	034703	412542	60000	4800	-1.15	4.50	0.6	-105	6	-130	17	-80	9	FJ98	Y
G 175-7	034828	511406	11000	5000	-1.76	4.00	1.0	-164	240	-515	400	-23	30	g-k	N
BD -4 680	035154	-034911	11000	5800	-1.89	4.00	1.0	-232	40	-185	58	77	58	g-k	N
GL 158	040315	351624	60000	4800	-1.79	5.00	1.0	-50	20	-185	4	26	1	FJ98	Y
G 39-36	044807	330936	60000	4200	-2.50	4.00	1.0	-14	8	-95	25	82	20	g-k	Y
G 81-41	045505	454405	60000	4550	-0.96	4.50	0.6	-8	4	-79	12	72	11	k	Y
BD -10 1085	050422	-100859	60000	5100	-0.50	4.50	0.8	-6	3	-70	4	-27	1	g-k	Y
BD +19 869	051253	194320	60000	4600	-1.01	4.50	0.5	0	1	-110	9	-5	1	k	Y
PLX 1219	052310	331130	30000	4600	-1.79	4.00	0.6	-240	15	-125	90	9	5	Yale	Y
HIP 27928	055434	-092334	60000	4200	-0.93	4.00	0.6	-70	3	25	3	45	4	Yale, HIP	Y
G 103-27	061648	295706	60000	5600	-0.95	4.50	0.6	0	5	-145	95	-118	80	K	N
G 101-34	062004	382044	60000	5000	-1.50	2.75	1.0	-336	374	-1425	4930	109	206	g ^d	N
HIP 30567	062530	484340	60000	4700	-0.60	4.75	0.6	8	4	-104	12	-46	5	k	Y
G 103-50	064008	282712	60000	4700	-2.20	4.50	0.4	-147	38	-71	3	15	24	AAM96, CLLA94	Y
AC +78 2199	065249	781230	60000	5000	-0.03	4.50	0.8	-120	19	-4	4	-114	23	g-k	Y
G 88-1	065828	185949	60000	4600	-0.84	5.00	1.0	2	3	-45	6	41	4	g-k	Y
BD -17 1716	065929	-171517	60000	4600	-0.08	4.50	0.4	-53	3	44	3	-25	3	k	Y
G 87-27	071008	371634	60000	5100	-0.61	3.50	0.9	-336	374	-1425	4930	109	206	CLLA94	Y
G 87-35	072044	292048	30000	5300	-0.53	4.50	0.8	-51	5	-123	21	50	6	g-k	Y
G 193-42	072222	503318	11000	5000	-0.94	4.25	1.0	-155	35	-275	140	37	10	g-k	N
G 193-49	072722	522612	30000	5000	-0.39	5.00	0.9	-72	26	35	14	-118	58	k	Y
G 90-4	073023	271618	11000	5600	-0.81	4.50	1.0	7	10	-150	120	-108	80	g-k	N
GSC 03790-02030	074534	565722	30000	5600	-1.16	4.00	0.7	-43	37	-84	50	-59	32	K	N
LP 208-60	082144	384713	11000	5800	-0.22	4.00	1.0	50	35	-280	255	34	37	g-k	N
G 115-1	082217	410424	30000	4800	-1.33	4.50	0.6	-75	3	-84	23	34	4	g-k	Y
LP 545-17	082540	051528	11000	5200	-1.01	4.75	1.0	52	105	-95	50	177	120	k-m	N
PLX 2019	082941	014448	30000	4600	-1.72	4.50	1.0	237	101	-334	104	-11	14	Yale	Y
PLX 2023.01	083000	-095401	30000	4400	-0.39	3.90	1.0	-12	1	4	1	10	1	Yale	Y
LP 209-14	083110	421348	11000	5400	-1.36	4.75	1.0	-155	16	-270	265	92	15	K	N
G 113-49	083512	032800	11000	5100	-0.83	4.00	1.0	95	120	-180	110	77	35	k-m	N
G 9-31	085245	223330	11000	5600	-1.00	4.25	1.0	-96	16	-150	45	-93	45	g-k	N
LP 36-78	085629	705814	60000	5150	0.25	4.50	0.7	-87	16	-56	23	31	2	k	Y
LP 786-61	090514	-183206	60000	5800	-0.76	4.00	0.8	63	23	16	3	47	20	g-k	N
ROSS 94	094722	261813	60000	4600	0.00	3.50	1.0	-24	2	-49	8	-18	4	k	Y
G 116-55	094819	340712	30000	5600	-2.10	4.00	1.0	-190	90	-165	130	-38	90	g-k	N
G 43-7	095013	050905	30000	4950	-1.02	4.50	0.8	113	19	-41	11	12	5	g-k	Y
G 161-84	095139	-035000	60000	4700	-1.31	4.75	0.4	-199	53	-96	17	131	15	CLLA94	Y

Table 1—Continued

Name	Ra ^a	Dec ^b	Resolving	T _{eff}	[Fe/H]	log g	ξ_t	U _{LSR}	σ_U	V _{LSR}	σ_V	W _{LSR}	σ_W	Luyten color	Molecular
	2000	2000	Power ^c	(K)		(cm s ⁻²)				(km/s)				or source	Features
LP 788-55	095433	-192100	60000	4700	-0.53	4.50	0.7	66	18	-4	6	-5	4	g-k	Y
WOLF 334	095744	323654	11000	5200	-1.71	4.50	1.0	-450	320	-375	335	-163	300	g-k	N
LP 315-12	095746	264522	60000	5100	-0.98	4.50	0.6	-122	15	-12	4	-14	11	g-k	Y
LP 429-17	100108	141842	60000	4750	-0.10	4.50	0.4	47	10	-47	13	-15	3	g-k	Y
BD +53 1395	101357	523024	60000	4500	-1.04	4.50	0.5	16	1	-79	3	12	1	k	Y
BD +12 2201	102220	120845	60000	4500	-0.23	4.50	0.9	-57	3	-17	1	-9	2	k	Y
LP 790-19	102607	-175843	60000	4300	-0.51	5.00	0.6	-111	7	-46	1	9	2	k	Y
BD -9 3104	103529	-102236	30000	4900	-0.60	4.50	0.8	130	32	-82	14	-9	9	k	Y
G 119-21	103725	285548	30000	4600	-0.48	4.50	0.6	-81	24	58	15	-42	10	k	Y
BD +31 2175	104016	304855	60000	4800	-0.17	4.75	1.1	-90	4	-35	2	-13	3	k	Y
PLX 2529.1	105203	-000938	60000	4800	-0.58	4.50	0.5	39	3	-6	3	-52	2	Yale, HIP	Y
G 119-45	105721	305948	30000	4800	0.14	4.50	0.7	-177	39	-85	22	-53	20	k	Y
G 58-36	110313	220024	30000	4600	-0.05	4.50	0.8	-115	35	-47	15	15	16	k	Y
G 253-46	110806	822454	60000	5200	-0.57	4.50	0.7	0	2	-35	6	56	7	g-k	Y
BD -10 3216	111111	-105703	60000	4500	-1.19	4.50	0.4	-122	3	-16	1	34	1	k-m	Y
HIP 55128	111712	172927	30000	4900	-0.38	4.50	0.9	-124	26	-55	14	-53	12	k	Y
LP 792-12	111858	-171548	60000	4800	-0.39	5.00	0.6	-77	14	-34	2	35	2	g-k	Y
BD +15 2325	112219	142644	60000	4800	-0.58	4.50	0.6	-125	12	-46	5	-9	5	k	Y
G 122-22	112400	453234	30000	4400	-1.36	4.50	0.9	63	15	-36	8	58	8	k	Y
G 163-80	112507	-055624	30000	4700	0.05	4.75	0.6	-114	23	-55	9	0	8	k	Y
ROSS 109	112715	593318	60000	5300	-1.53	4.00	0.6	-350	300	-645	700	217	140	K	N
LP 733-14	113829	-135006	60000	4700	-0.57	4.75	0.7	-15	3	-35	11	-41	10	g-k	Y
LP 734-54	115754	-094848	60000	4800	-0.66	4.50	0.4	-78	13	-15	4	5	2	g-k	Y
WOLF 1424	120019	203543	60000	4600	-1.19	4.50	0.4	-28	4	-120	25	-33	6	k	Y
LP 734-101	121655	-114906	60000	5500	0.06	4.50	0.7	-86	16	-36	10	-9	4	g-k	Y
G 13-29	121920	022642	30000	4800	-0.66	4.50	0.6	-6	4	-101	30	-8	15	g-k	Y
G 198-61	123523	374430	30000	5300	-0.12	4.50	1.0	176	36	-42	11	21	7	k	Y
G 149-9	124803	274548	30000	4900	-0.62	4.75	0.8	24	16	-315	150	-86	1	k	Y
HIP 62627	124956	711139	60000	4800	-0.24	5.00	1.2	-8	2	-93	3	-17	2	HIP	Y
G 149-18	125246	222700	30000	5000	-0.03	4.50	0.8	-214	44	-33	11	29	2	k	Y
W 453	125749	054554	30000	4600	-0.04	4.75	0.9	-71	17	-38	13	2	2	k	Y
HD 114095	130826	-071830	60000	4730	-0.71	2.40	1.3	65	43	-290	166	-49	32	AAM96	Y
BD +68 714	131021	672941	30000	4900	-0.05	4.50	1.0	-96	3	-70	2	7	1	K1	Y
ROSS 466	131514	-110112	30000	5000	-0.86	4.75	0.9	-71	17	-115	65	-103	30	g-k	Y
LHS 2715	131857	-030418	60000	4400	-1.56	4.00	0.6	-40	7	-120	7	115	1	Gizis97	Y
LP 172-89	132638	464851	30000	4800	-0.49	4.50	0.6	-124	28	-34	9	0	2	g-k	Y
LP 738-45	133814	-154714	30000	4400	-0.12	4.50	0.8	-81	17	-28	14	14	4	k	Y
G 255-44	135536	740012	60000	4900	-1.30	4.50	0.7	-107	17	-25	15	74	7	k	Y
G 65-22	140144	085517	60000	5000	-1.66	4.50	0.4	10	8	-225	76	-83	25	AAM96, CLLA94	Y
BD +30 2490	141124	300502	30000	4600	-0.18	4.75	0.6	-68	5	-88	8	35	3	k	Y
G 239-12	141853	731412	60000	5800	-2.62	4.00	1.0	74	25	-333	160	127	170	K	N
HIP 70152	142114	085816	60000	4800	-0.05	5.00	1.0	50	7	-36	5	-20	4	RWMRKM01	Y
HIP 70715	142745	235027	60000	4400	-0.11	4.50	0.6	-110	9	-66	6	-9	4	HIP	Y
LP 500-92	143413	123457	30000	4500	-0.38	4.50	0.9	-134	9	-37	4	10	5	k-m	Y
LP 175-8	143610	450859	30000	4900	-0.31	4.75	0.6	-123	25	-65	14	33	10	k	Y
HD 131287	145004	603936	30000	4800	-0.04	4.50	1.0	-4	1	-12	1	-13	1	k	Y
BD +23 2751	145342	232043	60000	4700	-0.51	4.50	0.6	-76	2	-62	2	20	1	k	Y
LP 502-8	150914	143123	30000	5000	-0.95	4.50	0.6	-80	17	-119	25	113	16	g-k	Y
HIP 74235	151013	-162246	60000	4900	-1.51	4.50	0.8	295	2	-505	18	-62	10	HIP	Y

Table 1—Continued

Name	Ra ^a	Dec ^b	Resolving	T _{eff}	[Fe/H]	log g	ξ_t	U _{LSR}	σ_U	V _{LSR}	σ_V	W _{LSR}	σ_W	Luyten color	Molecular
	2000	2000	Power ^c	(K)		(cm s ⁻²)				(km/s)				or source	Features
LP 802-56	151912	-210046	60000	5200	-0.88	4.50	0.6	-143	7	-130	32	-82	7	k	Y
BD +2 2944	151919	014555	30000	6000	-0.02	3.50	1.8	81	1	-8	1	-3	1	K4	N
ROSS 804	154529	-134918	30000	4600	-0.93	4.50	0.6	120	85	-285	360	-108	200	k	Y
G 225-50	161445	552549	30000	4600	-0.13	4.50	1.0	-60	13	-43	6	8	7	g-k	Y
LP 330-7	162306	320835	30000	4800	-0.22	4.50	0.6	74	16	4	6	10	6	g-k	Y
G 202-57	163337	511330	30000	4900	-1.05	4.50	0.6	180	82	-65	32	49	18	g-k	Y
G 17-25	163442	-041345	60000	4950	-1.46	4.50	0.3	11	11	-115	24	-29	6	CLLA94	Y
BD -14 4454	163614	-151012	30000	4600	-0.95	4.50	0.6	-111	2	-90	8	35	5	k	Y
LP 445-55	164258	192209	30000	5200	-1.80	4.00	0.6	118	24	38	8	84	13	g-k	N
LP 447-2	170805	175738	30000	5800	-1.79	4.00	0.9	-146	7	-272	26	46	18	g-k	N
LP 747-18	171644	-130112	30000	5800	-0.95	4.50	0.8	71	51	-322	327	41	29	g-k	N
LP 447-34	171708	180706	30000	4800	-0.35	4.50	0.9	53	18	-45	20	11	4	g-k	Y
G 19-25	172559	-024436	60000	4900	-2.01	4.50	0.4	116	86	-93	85	9	6	AAM96, CLLA94	Y
LP 808-11	173353	-165854	30000	5500	-1.23	4.00	0.7	-35	40	-225	200	-78	65	g-k	N
BD -8 4501	174728	-084648	30000	5800	-1.80	4.00	0.8	126	12	-45	16	-156	38	g-k	N
G 204-47	180929	471324	30000	5200	-0.88	4.50	1.0	170	126	-137	65	73	59	K	N
BD +5 3640	181222	052404	60000	4950	-1.34	4.50	0.4	61	13	-216	21	38	5	AAM96	Y
BD -17 5287	184057	-170954	30000	5800	-0.70	4.50	1.0	-60	25	-155	80	77	35	K	N
G 227-44	184143	583430	30000	5400	-1.45	4.50	1.0	-314	250	-210	11	-37	42	K	N
G 207-23	191613	370420	30000	5000	-1.90	4.00	0.6	-175	6	-319	1	41	15	K	N
HD 181007	191928	-202540	60000	4700	-1.93	1.50	1.6	65	43	-290	166	-49	32	None ^e	N
LP 752-18	192547	-110954	30000	5800	-1.00	4.00	1.0	60	25	-75	40	17	10	g-k	N
LP 753-29	195004	-131912	30000	5400	-1.88	4.00	1.0	-65	26	-155	76	77	35	g-k	N
G 23-14	195150	053646	60000	5000	-1.51	3.00	1.0	116	86	-93	85	9	6	g ^f	N
LP 575-39	202326	055050	30000	4700	-0.42	4.50	0.6	-90	13	-11	7	-19	12	g-k	Y
LP 575-40	202356	052444	30000	4800	-0.10	4.50	1.0	-110	25	20	15	-29	18	k-m	Y
LP 815-43	203814	-202554	30000	6400	-2.70	4.00	1.0	140	140	-290	265	36	27	g-k	N
HD 200968	210710	-135523	30000	5000	-0.16	4.50	1.2	-52	1	-14	1	2	1	KF99	Y
BD +22 4567	221031	224749	30000	4800	-0.33	4.75	1.1	75	4	-5	1	70	3	k	Y
BD +30 4633	221206	313341	60000	4600	-0.01	4.50	0.8	92	6	-35	1	-6	2	k-m	Y
HIP 109801	221424	-084442	30000	4600	-1.70	4.50	1.0	130	50	-225	75	-43	25	HIP	Y
ROSS 237	225330	274518	30000	4900	-1.49	4.25	0.9	-118	25	-315	16	8	25	k	Y
ROSS 242	230849	270054	60000	4700	-0.94	4.50	0.4	50	14	-130	16	-67	30	k	Y
LP 702-79	232332	-060012	30000	4800	0.05	4.50	1.1	-150	60	-70	25	1	22	g-k	Y
HIP 115664	232556	291141	30000	4600	-0.90	5.00	0.6	-85	16	-215	20	-78	25	HIP	Y
BD +28 4634	234510	293343	60000	5000	-0.30	4.50	0.7	-138	3	6	2	-60	1	k	Y

^ahhmmss^bddmmss^cHET data taken with R=11000, 2.7m data taken with R=30000,60000^dStar chosen from Carney et al. (1994) and Alonso et al. (1996a) but also in NLTT catalogue^eStar chosen from Alonso et al. (1996a) not in NLTT catalogue^fStar chosen from Carney et al. (1994) but also in NLTT catalogue

References. — (AAM96) = Alonso et al. (1996a); (CLLA94) = Carney et al. (1994); (FJ98) = Fuchs & Jahreiß (1998); (KF99) = Kotoneva & Flynn (1999, private communication) (Gizis97) = Gizis (1997); (HIP) = HIPPARCOS color-magnitude diagram; (RWMRKM01) = Reid et al. (2001); (Yale) = Yale color-magnitude diagram

Table 2. Comparison with literature

Star	This study		Literature		Source
	T_{eff}	[Fe/H]	T_{eff}	[Fe/H]	
PLX 5805	4600	-1.72	...	-1.28	SPC93
LP 646-24	4800	-1.01	...	-0.24	RN91
G 72-34	4800	-2.23	4689	-2.21	CLLA94
BD +8 335	5100	-0.98	5224	-0.80	CLLA94
	5100	-0.98	...	-0.61	Eggen98
BD +4 415	4750	-0.63	4756	-0.41	AAM96
	4750	-0.63	...	-0.43	Reid01
	4750	-0.63	...	-0.39	Eggen98
G 78-26	4200	-1.01	...	-1.20	Gizis97
GJ 1064A	5000	-1.15	5049	-1.02	CLLA94
	5000	-1.15	5200	-1.05	TL99
	5000	-1.15	...	-0.85	TI99
	5000	-1.15	5050	-0.80	Fulbright00
GJ 1064B	4800	-1.15	4950	-1.11	TL99
	4800	-1.15	4816	-1.05	CLLA94
	4800	-1.15	...	-0.86	TI99
BD -4 680	5800	-1.89	5866	-2.07	AAM96
GL 158	4800	-1.79	4762	-1.73	CLLA94
	4800	-1.79	4700	-1.70	Fulbright00
	4800	-1.79	...	-1.67	TI99
	4800	-1.79	4775	-1.81	TL99
BD +19 869	4600	-1.01	...	-0.55	Eggen87
	4600	-1.01	...	-0.60	Norris86
PLX 1219	4600	-1.79	4935	-1.02	CLLA94
	4600	-1.79	...	-2.00	Morrison01
	4600	-1.79	4739	-1.61	AAM96
G 103-27	5600	-0.95	5449	-0.93	SPC93
G 101-34	5000	-1.50	5289	-1.83	AAM96 ^a
	5000	-1.50	4955	-1.88	CLLA94
G 103-50	4700	-2.20	4642	-2.00	CLLA94
	4700	-2.20	4713	-3.00	AAM96
G 88-1	4600	-0.84	...	-0.13	RN91
	4600	-0.84	...	-0.40	SPC93
G 87-27	5100	-0.61	4905	-1.45	CLLA94 ^a
	5100	-0.61	5090	-0.42	GL00
G 87-35	5300	-0.53	5222	-0.48	CLLA94
	5300	-0.53	...	-0.55	Eggen98
G 90-4	5600	-0.81	5513	-0.86	CLLA94
PLX 2019	4600	-1.72	4740	-1.13	AAM96
G 113-49	5100	-0.83	5194	-0.78	CLLA94
G 9-31	5600	-1.00	5546	-0.94	CLLA94
	5600	-1.00	5569	-0.85	AAM96
G 43-7	4950	-1.02	5009	-0.68	CLLA94
	4950	-1.02	...	-0.68	Reid01
G 161-84	4700	-1.31	4650	-1.71	CLLA94
	4700	-1.31	...	-1.57	Beers99
WOLF 334	5200	-1.71	5084	-1.82	CLLA94
LP 792-12	4800	-0.39	...	0.18	RN91
G 13-29	4800	-0.66	...	0.04	RN91
HD 114095	4730	-0.71	4650	-0.60	Fulbright00
	4730	-0.71	4541	-1.58	AAM96 ^a
	4730	-0.71	4680	-0.35	Clementini99
BD +68 714	4900	-0.05	...	-0.17	Eggen98
LHS 2715	4400	-1.56	4480	-0.22	AAM96
	4400	-1.56	...	-1.20	Gizis97
G 65-22	5000	-1.66	4971	-1.75	AAM96
	5000	-1.66	4982	-1.72	CLLA94
G 239-12	5800	-2.62	6025	-2.56	CLLA94
HIP 70152	4800	-0.05	...	0.22	RN91
	4800	-0.05	...	-0.03	Reid01
HIP 74235	4900	-1.51	5040	-1.54	TL99
	4900	-1.51	...	-1.31	TI99
	4900	-1.51	4957	-1.57	CLLA94
	4900	-1.51	4974	-1.52	AAM96
	4900	-1.51	4850	-1.40	Fulbright00

Table 2—Continued

Star	This study		Literature		Source
	T _{eff}	[Fe/H]	T _{eff}	[Fe/H]	
BD +2 2944	6000	-0.02	6036	-0.13	AAM99
G 17-25	4950	-1.46	5175	-1.10	Fulbright00
	4950	-1.46	...	-1.08	TI99
	4950	-1.46	5180	-1.35	TL99
	4950	-1.46	4966	-1.34	AAM96
	4950	-1.46	4959	-1.54	CLLA94
LP 445-55	5200	-1.80	...	-1.55	RN91
LP 447-2	5800	-1.79	...	-1.74	RN91
LP 747-18	5800	-0.95	...	-1.22	RN91
G 19-25	4900	-2.01	...	-2.03	RN91
	4900	-2.01	4938	-1.88	CLLA94
	4900	-2.01	4977	-1.52	AAM96
BD -8 4501	5800	-1.80	5657	-1.99	CLLA94
	5800	-1.80	5682	-2.79	AAM96
G 204-47	5200	-0.88	5249	-0.86	CLLA94
BD +5 3640	4950	-1.34	5080	-1.27	TL99
	4950	-1.34	5012	-1.36	CLLA94
	4950	-1.34	4980	-1.44	AAM96
	4950	-1.34	...	-0.37	RN91
G 227-44	5400	-1.45	5501	-1.41	CLLA94
G 207-23	5000	-1.90	...	-1.00	RWMRKM01
HD 181007	4700	-1.93	4582	-2.27	AAM96 ^a
LP 752-18	5800	-1.00	...	-1.73	RN91
LP 753-29	5400	-1.88	...	-2.10	RN91
G 23-14	5000	-1.51	5100	-1.40	Carney97
	5000	-1.51	4893	-2.20	CLLA94
LP 815-43	6400	-2.70	6379	-2.64	TI99
	6400	-2.70	6500	-2.95	PMBH00
BD +22 4567	4800	-0.33	...	-0.22	Eggen98
HIP 109801	4600	-1.70	4661	-1.43	CLLA94
	4600	-1.70	...	-0.92	RN91
ROSS 237	4900	-1.49	4947	-1.38	CLLA94
BD +28 4634	5000	-0.30	...	-0.15	SPC93

^aGiant star assumed to be a dwarf

Note. — The following references were based on spectroscopic data: Beers99, CLLA94, Carney97, Clementini99, Fulbright00, Gizis97, Primas00, RN91, TI99, TL99.

References. — (AAM96) = Alonso et al. (1996a); (AAM99) = Alonso et al. (1999a); (Beers99) = Beers et al. (1999); (CLLA94) = Carney et al. (1994); (Carney97) = Carney et al. (1997); (Clementini99) = Clementini et al. (1999); (Eggen87) = Eggen (1987); (Eggen98) = Eggen (1998); (Fulbright00) = Fulbright (2000); (GL00) = Gay & Lambert (2000); (Gizis97) = Gizis (1997); (Morrison01) = Morrison et al. (2001); (Norris86) = Norris (1986); (Primas00) = Primas et al. (2000); (Reid01) = Reid et al. (2001); (RN91) = Ryan & Norris (1991); (SPC93) = Schuster et al. (1993); (TI99) = Thévenin & Idiart (1999); (TL99) = Tomkin & Lambert (1999)

Table 3. Comparison of T_{eff} 's derived in this study with T_{eff} 's calculated from the Alonso et al. (1996b, 1999b) relations for $T_{\text{eff}}:[Fe/H]:color$ using photometry found in the literature.

Star	T_{eff} (K) This study	[Fe/H]	Color Index	T_{eff} (K) from color relation
PLX 5805	4600	-1.72	B-V	4764
	4600	-1.72	J-K	4464
BD -14 142	6200	-1.21	B-V	5113
	6200	-1.21	V-K	6327
	6200	-1.21	J-K	5860
LP 646-24	4800	-1.01	V-K	4666
	4800	-1.01	J-K	4665
G 72-34	4800	-2.23	J-K	4654
BD -4 290	4900	-0.01	B-V	4544
BD +8 335	5100	-0.98	B-V	4485
HIP 10412	4600	-0.40	B-V	4491
Cool 340	4800	-0.40	B-V	4581
	4800	-0.40	V-K	4158
	4800	-0.40	J-K	4544
BD +4 415	4750	-0.63	AAM96 ^a	4756
GJ 1064A	5000	-1.15	B-V	4717
	5000	-1.15	V-K	4963
	5000	-1.15	J-K	4978
GJ 1064B	4800	-1.15	B-V	4692
	4800	-1.15	V-K	4591
	4800	-1.15	J-K	5791
BD -4 680	5800	-1.89	AAM96 ^a	5866
G 39-36	4200	-2.50	J-K	4400
G 81-41	4550	-0.96	B-V	4357
BD -10 1085	5100	-0.50	B-V	4676
	5100	-0.50	V-K	4982
	5100	-0.50	J-K	4978
BD +19 869	4600	-1.01	V-K	4245
	4600	-1.01	J-K	4287
PLX 1219	4600	-1.79	AAM96 ^a	4739
HIP 27928	4200	-0.93	J-K	4082
G 103-27	5600	-0.95	B-V	5580
	5600	-0.95	V-K	5118
	5600	-0.95	J-K	5153
G 101-34	5000	-1.50	B-V	4724 ^b
G 103-50	4700	-2.20	AAM96 ^a	4713
HIP 30567	4700	-0.60	B-V	4701
AC +78 2199	5000	-0.03	B-V	4408
	5000	-0.03	V-K	5036
	5000	-0.03	J-K	5026
G 88-1	4600	-0.84	B-V	4700
	4600	-0.84	V-K	4394
	4600	-0.84	J-K	4482
BD -17 1716	4600	-0.08	B-V	4438
	4600	-0.08	V-K	4397
	4600	-0.08	J-K	4301
G 87-27	5100	-0.61	B-V	4697 ^b
G 87-35	5300	-0.53	B-V	6191
	5300	-0.53	V-K	5055
	5300	-0.53	J-K	5242
G 193-49	5000	-0.39	B-V	4538
G 90-4	5600	-0.81	b-y,c1	5473
LP 208-60	5800	-0.22	J-K	5121
G 115-1	4800	-1.33	B-V	4407
	4800	-1.33	V-K	4884
	4800	-1.33	J-K	4784
PLX 2019	4600	-1.72	AAM96 ^a	4740
LP 209-14	5400	-1.36	J-K	5701
G 9-31	5600	-1.00	AAM96 ^a	5569
LP 36-78	5150	0.25	B-V	4786
LP 786-61	5800	-0.76	B-V	5557
ROSS 94	4600	0.00	B-V	3652 ^b
	4600	0.00	V-K	3744 ^b
	4600	0.00	J-K	4207 ^b
G 116-55	5600	-2.10	B-V	5513
	5600	-2.10	V-K	5740

Table 3—Continued

Star	T _{eff} (K) This study	[Fe/H]	Color Index	T _{eff} (K) from color relation
	5600	-2.10	J-K	5486
G 43-7	4950	-1.02	B-V	6627
G 161-84	4700	-1.31	J-K	4471
LP 788-55	4700	-0.53	J-K	4612
WOLF 334	5200	-1.71	J-K	5455
LP 315-12	5100	-0.98	V-K	4582
	5100	-0.98	J-K	4756
LP 429-17	4750	-0.10	B-V	4658
	4750	-0.10	V-K	4563
	4750	-0.10	J-K	4604
BD +12 2201	4500	-0.23	V-K	4245
	4500	-0.23	J-K	4095
LP 790-19	4300	-0.51	J-K	4119
BD -9 3104	4900	-0.60	B-V	4853
	4900	-0.60	V-K	4785
	4900	-0.60	J-K	4884
G 119-21	4600	-0.48	B-V	4126
	4600	-0.48	V-K	4304
	4600	-0.48	J-K	4604
BD +31 2175	4800	-0.17	B-V	4502
	4800	-0.17	V-K	4826
	4800	-0.17	J-K	4673
PLX 2529.1	4800	-0.58	B-V	4610
	4800	-0.58	V-K	4653
	4800	-0.58	J-K	4650
G 119-45	4800	0.14	J-K	4537
G 58-36	4600	-0.05	J-K	4540
G 253-46	5200	-0.57	V-K	5300
	5200	-0.57	J-K	5394
HIP 55128	4900	-0.38	B-V	4609
	4900	-0.38	V-K	4685
	4900	-0.38	J-K	4728
LP 792-12	4800	-0.39	B-V	4261
G 122-22	4400	-1.36	V-K	4465
G 163-80	4700	0.05	J-K	4658
ROSS 109	5300	-1.53	B-V	5610
LP 733-14	4700	-0.57	V-K	4534
	4700	-0.57	J-K	4407
WOLF 1424	4600	-1.19	V-K	4491
	4600	-1.19	J-K	4358
LP 734-101	5500	0.06	B-V	5548
G 13-29	4800	-0.66	V-K	4684
	4800	-0.66	J-K	4338
G 198-61	5300	-0.12	B-V	5147
	5300	-0.12	V-K	5139
	5300	-0.12	J-K	5460
G 149-9	4900	-0.62	V-K	4731
	4900	-0.62	J-K	4631
HIP 62627	4800	-0.24	B-V	4295
G 149-18	5000	-0.03	B-V	4206
	5000	-0.03	V-K	4643
	5000	-0.03	J-K	4650
HD 114095	4730	-0.71	B-V	4422 ^b
	4730	-0.71	V-K	4510 ^b
	4730	-0.71	J-K	4565 ^b
	4730	-0.71	AAM96 ^a	4541 ^c
BD +68 714	4900	-0.05	B-V	4424
	4900	-0.05	V-K	4797
	4900	-0.05	J-K	4952
ROSS 466	5000	-0.86	J-K	4784
LHS 2715	4400	-1.56	AAM96 ^a	4480
G 65-22	5000	-1.66	AAM96 ^a	4971
LP 172-89	4800	-0.49	B-V	4384
LP 738-45	4400	-0.12	J-K	4264
G 255-44	4900	-1.30	B-V	4901
	4900	-1.30	V-K	4686

Table 3—Continued

Star	T _{eff} (K) This study	[Fe/H]	Color Index	T _{eff} (K) from color relation
	4900	-1.30	J-K	4813
BD +30 2490	4600	-0.18	B-V	4005
	4600	-0.18	V-K	4185
	4600	-0.18	J-K	4442
G 239-12	5800	-2.62	B-V	5460
HIP 70152	4800	-0.05	B-V	4446
LP 500-92	4500	-0.38	B-V	3765
	4500	-0.38	V-K	4184
	4500	-0.38	J-K	4294
LP 175-8	4900	-0.31	B-V	4490
	4900	-0.31	V-K	4678
	4900	-0.31	J-K	4842
HD 131287	4800	-0.04	B-V	5715
LP 502-8	5000	-0.95	B-V	4559
HIP 74235	4900	-1.51	AAM96 ^a	4971
BD +2 2944	6000	-0.02	AAM99 ^a	5856 ^b
G 225-50	4600	-0.13	B-V	3964
	4600	-0.13	V-K	4637
	4600	-0.13	J-K	4341
LP 330-7	4800	-0.22	B-V	4087
	4800	-0.22	V-K	4743
	4800	-0.22	J-K	4708
G 202-57	4900	-1.05	B-V	4917
	4900	-1.05	V-K	4617
	4900	-1.05	J-K	4612
G 17-25	4950	-1.46	AAM96 ^a	4966
BD -14 4454	4600	-0.95	V-K	4474
	4600	-0.95	J-K	4334
LP 445-55	5200	-1.80	V-K	5807
	5200	-1.80	J-K	5420
LP 447-2	5800	-1.79	B-V	5404
	5800	-1.79	V-K	5611
	5800	-1.79	J-K	5491
LP 747-18	5800	-0.95	B-V	5380
	5800	-0.95	V-K	4827
	5800	-0.95	J-K	5103
LP 447-34	4800	-0.35	B-V	5136
	4800	-0.35	V-K	4699
	4800	-0.35	J-K	4760
G 19-25	4900	-2.01	AAM96 ^a	4977
BD -8 4501	5800	-1.80	AAM96 ^a	5682
G 204-47	5200	-0.88	B-V	4923
	5200	-0.88	V-K	5044
	5200	-0.88	J-K	5394
BD +5 3640	4950	-1.34	AAM96 ^a	4980
BD -17 5287	5800	-0.70	B-V	4584
G 227-44	5400	-1.45	B-V	4932
G 207-23	5000	-1.90	B-V	4798
	5000	-1.90	V-K	5279
	5000	-1.90	J-K	5176
HD 181007	4700	-1.93	B-V	4659 ^b
	4700	-1.93	V-K	4676 ^b
	4700	-1.93	J-K	4887 ^b
LP 752-18	5800	-1.00	B-V	6055
G 23-14	5000	-1.51	b-y	4958 ^b
LP 575-39	4700	-0.42	B-V	4100
LP 575-40	4800	-0.10	B-V	4349
LP 815-43	6400	-2.70	B-V	5193
	6400	-2.70	V-K	6509
	6400	-2.70	J-K	5973
HD 200968	5000	-0.16	B-V	4550
BD +22 4567	4800	-0.33	B-V	4315
BD +30 4633	4600	-0.01	B-V	3889
HIP 109801	4600	-1.70	J-K	4593
ROSS 237	4900	-1.49	B-V	5410
	4900	-1.49	V-K	4590

Table 3—Continued

Star	T _{eff} (K) This study	[Fe/H]	Color Index	T _{eff} (K) from color relation
ROSS 242	4900	-1.49	J-K	4879
	4700	-0.94	B-V	4680
	4700	-0.94	V-K	4682
	4700	-0.94	J-K	4646
LP 702-79	4800	0.05	V-K	4605
	4800	0.05	J-K	4331
HIP 115664	4600	-0.90	V-K	4370
	4600	-0.90	J-K	4411
BD +28 4634	5000	-0.30	B-V	4515
	5000	-0.30	V-K	4825
	5000	-0.30	J-K	4712

^aT_{eff}'s determined from direct application of the IRFM

^bGiant star; T_{eff} derived from Alonso et al. (1999b) relation

^cGiant star assumed to be a dwarf

Note. — Photometry from the following sources. B and V taken from the Guide Star Catalog II. J and K taken from the Two Micron All Sky Survey (2MASS) Skrutskie et al. (1997). b-y and c1 taken from Schuster et al. (1993).

References. — (AAM96) = Alonso et al. (1996a); (AAM99) = Alonso et al. (1999a)